

Hubble Space Telescope Observations of Binary Very-Low-Mass Stars and Brown Dwarfs

John E. Gizis¹

Department of Physics and Astronomy, University of Delaware, Newark, DE 19716

`gizis@udel.edu`

I. Neill Reid

Space Telescope Science Institute, 3700 San Marin Drive, Baltimore, MD 21218

Gillian R. Knapp

Department of Astrophysical Sciences, Princeton University, Princeton, NJ 08544-1001

James Liebert

Steward Observatory, University of Arizona, Tucson, AZ 85721

J. Davy Kirkpatrick

Infrared Processing and Analysis Center, 100-22, California Institute of Technology, Pasadena, CA 91125

David W. Koerner

Department of Physics and Astronomy, Northern Arizona University, NAU Box 6010, Building 19, Flagstaff, AZ 86011-6010

Adam J. Burgasser²

UCLA Division of Astronomy & Astrophysics, 8965 Math Science Bldg., 405 Hilgard Ave., Los Angeles, CA, 90095-1562

ABSTRACT

We present analysis of Hubble Space Telescope images of 82 nearby field late-M and L dwarfs. We resolve 13 of these systems into double M/L dwarf systems and identify

¹Visiting Astronomer, Kitt Peak National Observatory, National Optical Astronomy Observatories, which is operated by the Association of Universities for Research in Astronomy, Inc. (AURA) under cooperative agreement with the National Science Foundation.

²Hubble Fellow

an additional possible binary. Combined with previous observations of 20 L dwarfs, we derive an observed binary fraction for ultracool dwarfs of $17^{+4}_{-3}\%$, where the statistics included systems with separations in the range 1.6-16 A.U. We argue that accounting for biases and incompleteness leads to an estimated binary fraction $15 \pm 5\%$ in the range 1.6-16 A.U. No systems wider than 16 A.U. are seen, implying that the wide companion frequency is less than 1.7%; the distribution of orbital separation is peaked at $\sim 2 - 4$ A.U. and differs greatly from the G dwarf binary distribution. Indirect evidence suggests that the binary fraction is $\sim 5 \pm 3\%$ for separations less than 1.6 A.U. We find no evidence for differences in the binary fraction between stellar late-M and L dwarfs and substellar L dwarfs. We note, however, that the widest (> 10 A.U.) systems in our sample are all of earlier (M8-L0) spectral type; a larger sample is needed to determine if this is a real effect. One system with a spectral type of L7 has a secondary that is fainter in the HST F814W filter but brighter in F1042M; we argue that this secondary is an early-T dwarf.

Subject headings: binaries: general — stars: low-mass, brown dwarfs

1. Introduction

The Deep Near-Infrared Survey (Epchtein 1997; Delfosse et al. 1997, DENIS), Two Micron All-Sky Survey (Skrutskie et al. 1997; Cutri et al. 2000, 2MASS), and Sloan Digital Sky Survey (York et al. 2000, SDSS) have enabled the discovery of numerous cool dwarfs and led to the definition of new spectral types L (Kirkpatrick et al. 1999; Martín et al. 1999a) and T (Burgasser et al. 2002a; Geballe et al. 2002). With effective temperatures lower than $\sim 2000\text{K}$ and $\sim 1300\text{K}$ respectively, these objects span the range between the lowest mass stars and sub-stellar mass brown dwarfs. The latest M dwarfs (M8/M9) and the earlier-type L dwarfs represent a mix of brown dwarfs which are still cooling and long-lived hydrogen-burning stars which have reached a stable state (Reid et al. 1999). Theoretical models indicate that the overwhelming majority of L dwarfs with temperature below $\sim 1750\text{K}$ (spectral types $\approx \text{L5}$ and later; Kirkpatrick et al. 1999, Gizis et al. 2000), and all T dwarfs, are expected to be brown dwarfs.

The large samples now available of these objects allow investigation of their statistical characteristics. One goal of such studies is the establishment of empirical constraints on the formation mechanism(s) of low mass dwarfs, particularly searching for potential differences in the properties of the lowest mass stars and brown dwarfs. It is clear that the properties of binary brown dwarfs are important in this context. A comprehensive theory of star formation must account not only for the stellar/substellar initial mass function (IMF), but also the frequency and orbital distributions of binary systems. Although brown dwarfs are extremely rare as close companions to GFK stars (Halbwachs et al. 2000; Marcy & Butler 2000) a number of studies have detected double brown dwarf systems (Martín et al. 1999b; Koerner et al. 1999; Reid et al. 2000; Close et al. 2002a; Bur-

gasser et al. 2003), and even doubles that orbit more massive stars (Martín et al. 2000a; Potter et al. 2002).³

The present paper presents analysis of the largest sample of high-resolution images of low mass dwarfs assembled so far. We have used the Planetary Camera of WFPC2 on the Hubble Space Telescope WFPC2 to image a sample of 82 late-type M and L dwarfs. Details of those observations are presented in Section 2. Comments on individual systems are given in Section 3. We have combined our present observations with data for twenty L dwarfs analyzed by Reid et al. (2000), and use those data to set constraints on the binary fraction in low mass dwarfs. That analysis is presented in Section 4, and our conclusions summarized in the final section.

2. Data

Targets for Snapshot WFPC2 imaging with HST were chosen from the list of 2MASS-selected late-M dwarfs (Gizis et al. 2000) and L dwarfs (Kirkpatrick et al. 1999, 2000), and from late-M and L dwarfs identified from the initial regions covered by SDSS (Fan et al. 2000; Schneider et al. 2002; Hawley et al. 2002). A few 2MASS and SDSS objects included have not yet been described in publications. Near-infrared discovery spectra for 2M0033-1521, 2M0326-2102 and 2M2242+2542 are shown in Figure 1. These data were obtained using the infrared Cryogenic Spectrometer (CRSP) on the Kitt Peak 4-meter telescope during observing runs in February and November 1999. The February data were obtained using Grating 2. The setup provided L-band spectra from K-band spectra from 1.85 to 2.55 μm . In November, we switched to the slightly higher resolution Grating 4, which yielded spectra over the range 1.90 to 2.55 μm at K-band or 1.05 to 3.75 μm at J-band. The resolution, as measured by the FWHM of arc lines, was 55 Å and 110 Å at J and K respectively for Grating 4. Data from this run are discussed in Schweitzer et al. (2002). Observations were obtained in the standard way described in the CRSP User’s Manual (Joyce 1999). Each object was stepped along the slit to obtain observations at five positions. When necessary, this cycle was repeated to obtain higher signal-to-noise. Typical individual exposures were 20 seconds at K-band. Spectra were extracted using IRAF. Both the HST photometry described below and comparison of these near-IR spectra to 25 optically classified L dwarfs observed with the same setup indicates that they are L dwarfs; however, optical spectra are needed to obtain accurate spectral types.

Each object on the HST program was observed at least once using the F814W and F1042M filters. For the brighter targets, a second F814W observation was possible in the timeframe allowed by the Snapshot visit. Exposure times in F814W ranged from 40 to 400 seconds while exposure times in F1042M were always 500 seconds. A total of 82 dwarfs were observed (Table 1). In general, Gizis & Reid (1995) found that we are able to resolve systems with magnitude differences of 0, 1, 3

³After the initial submission of this paper, Close et al. (2003) reported the independent discovery that 2M1127+74AB and 2M1311+80AB are doubles.

and 5 at separations of 0.09, 0.14, 0.23, and 0.31 arcseconds respectively using WFPC2. It should be noted, however, than in the case of the fainter L dwarfs, necessarily observed with lower signal-to-noise, that only companions within 3 magnitudes of the primary are detectable. We searched only the PC chip for companions. Coverage for wider companions is provided by the 2MASS data, which would have detected any companions ($K_s < 14.5$) wider than a few arcseconds.

Aperture photometry was measured and zero-pointed for each target according to the precepts of Holtzman et al. (1995). However, since we usually have only single frames in each filter, we found that the large aperture magnitudes were often contaminated by cosmic rays. Furthermore, the aperture magnitudes cannot be used for close doubles. DAOPHOT/IRAF was therefore used to measure magnitudes by fitting a PSF model, which was built using images of bright, single stars from this program. The PSF-subtracted images were an effective way of verifying that no secondaries were missed in the apparently single stars. The normalization between the DAOPHOT magnitudes and the aperture magnitudes was determined by taking the median value for high signal-to-noise, isolated stars. Finally, we corrected for charge transfer efficiency (CTE) losses using Dolphin (2000)’s Y-CTE corrections and X-CTE corrections. The resulting magnitudes for each system are given in Table 1. This table includes 2MASS Version-2 Working Database (corresponding to the Second Incremental Data Release system) JHK_s magnitudes for all systems. None of the binaries were resolved by 2MASS, so the magnitudes listed reflect the combined photometry. Table 1 lists composite WFPC2 photometry for those systems. Magnitudes and separations for the individual components in the resolved systems are given in Table 2. Z-band photometry of 2M0335+2342, 2M1711+2232, 2M2208+2921, and 2M2306-0502 was not possible due to cosmic ray hits. It is possible that low-level cosmic ray hits might also affect other targets, especially the Z band.

Two of the systems (2M0345+2540, 2M2224-0158) in our sample have trigonometric parallax measurements (Dahn et al. 2002). For the remainder, we have estimated photometric parallaxes. Unfortunately, the $I - Z$ WFPC2-based colors are not of high enough accuracy to directly estimate photometric distances for each component. Our distances are therefore based on combining the observed I magnitude for the primary, the Reid et al. (2000) $I_C - I_{814}$ relation, and the Dahn et al. (2002) $I_C - J, M_J$ relation.⁴ We assume the system $I - J$ color corresponds to the primary’s $I - J$ color in order to use the Dahn et al. (2002) relation; we tested differing assumptions such as using the observed system spectral type and got consistent answers. The transformation from separation in arcseconds to A.U. is uncertain by $\sim 30\%$; trigonometric parallaxes are needed. These errors have no significance for the interpretation of the data.

Color-color diagrams combining the HST and 2MASS photometry are shown in Figures 2 and 3. T dwarfs with WFPC2 photometry by Burgasser et al. (2003) are also shown. The gap between the latest L dwarfs of our program and the T5 and later dwarfs is obvious and is due to the lack of WFPC2 observations of early-T dwarfs. Our reddest object in $I-Z$ is the latest L dwarf, the L8

⁴We have adopted this procedure because our new sample adds only one object with an I_C measure, and because many of the parallax stars lack WFPC2 measurements.

2M0328+2302.

3. Individual Systems

2M1239+5515: This system shows lithium absorption according to Kirkpatrick et al. (2000), the only resolved binary in this study to do so. As such, both components of the system are below the lithium-burning limit of $0.055 - 0.060 M_{\odot}$ (Chabrier & Baraffe 1997; Burrows et al. 2001).

2M1728+3948: This system was classified L7 by Kirkpatrick et al. (2000). Remarkably, component A is brighter by 0.3 magnitudes at I but *fainter* by 0.3 magnitudes at Z than component B. Component B, with I-Z color of 3.1, is redder than the latest-type L dwarfs (Figures 2 and 3), suggesting that it lies in the L/T transition régime. This behaviour invites comparison with observations of late-L and early-T dwarfs at $1.25 \mu\text{m}$, where the latter dwarfs are observed to be brighter by up to 1 magnitude (Dahn et al. 2002). This has been interpreted as evidence for cloud disruption at the atmospheric temperatures spanned by the L/T transition (Burgasser et al. 2002b). Our observations provide the first evidence that the effect extends down to 1 micron. The primary in this system is cool enough that both components are brown dwarfs, with at least the primary above the lithium-burning limit.

2M1017+1308, 2M1430+2915: These two objects were identified as L dwarfs by J. Wilson (2002, priv. comm.) using near-IR spectra. The HST colors confirm the cool dwarf classification. Given the spectral types, all of the components could be either stars or brown dwarfs.

2M1449+2355: The limits on lithium for this distant L0 dwarf (Kirkpatrick et al. 2000) system do not rule out its presence. Both components may be stars or brown dwarfs. At 13 A.U., this is the most widely-separated L dwarf double currently known.

2M1600+1708: LRIS observations of this L1.5 dwarf (Kirkpatrick et al. 2000) system show no evidence for lithium absorption, indicating $M > 0.055 - 0.060 M_{\odot}$. The HST image is better modelled by two PSFs separated by 0.056 arcseconds than by a single PSF; we regard this as only a *candidate* double. Additional observations are required to demonstrate that this truly is a binary system.

2M2101+1756: This L7.5 dwarf (Kirkpatrick et al. 2000) system is late enough that both components must be brown dwarfs. The secondary is probably an L8 or slightly later dwarf.

2M1426+1557, 2M2140+1625, 2M2206-2047, 2M2331-0406: These late-M dwarf systems have primaries that are more likely to be stars than brown dwarfs; if any of these systems are young then the primary may be a brown dwarf. All four systems were discovered by Close et al. (2002a,b) using Gemini Adaptive Optics J, H and K-band imaging, and to whose discussion of individual systems we refer the reader. The *IZ* photometry presented here is consistent with the spectral types expected from the JHK photometry. 2M2206-2047 has a low tangential velocity (10 km/s), which is consistent with youth; since many old stars have small velocities, it does not prove

youth. Reid et al. (2002) have published a high resolution spectrum of 2M2206-2047 which shows rapid rotation and no lithium; thus, the possibility that the primary is a high-mass brown dwarf remains viable, but the mass must be above the lithium-burning limit.

2M1127+7411, 2M1311+8032, 2M2147+1431: These late-M dwarf systems have primaries that are more likely to be stars than brown dwarfs; if any of these systems are young then the primary may be a brown dwarf. Using our distance estimate and the Gizis et al. (2000) proper motion, the tangential velocity of 2M1127+7411 is only 6 km/s; this suggests youth but is not definitive. Our measurements of 2M1127+74AB and 2M1311+80AB are consistent with those reported by Close et al. (2003).

2MASSW0335+2342: This M8.5 dwarf shows lithium absorption and therefore is young brown dwarf (Reid et al. 2002). Our observations show no evidence for binarity.

4. The Binary Fraction

The most notable characteristic of the data is the lack of wide systems. Even for our faintest, most distant targets, we would have resolved companions with separations > 20 A.U., yet none are seen. Combining our sample with Reid et al. (2000), we find that 0 out of 102 late-M and L dwarfs have such companions with $\Delta I < 3$ (and in many cases, $\Delta I < 5$). The 1-sigma upper limit on the > 20 A.U. companion frequency is then 1.7%.

Thirteen of the 82 systems in the present sample are resolved as binaries with component separations exceeding our detection thresholds, all in the range 3-16 A.U. We have combined these data with the sample of twenty L dwarfs from Reid et al. (2000). Those dwarfs were also identified by color selection from the 2MASS database, and four are clearly resolved as binary systems with $\Delta > 0.06$ arcsec. This total observed binary fraction of $17^{+4}_{-3}\%$, where the uncertainties are calculated according to Burgasser et al. (2003), must be corrected to obtain a true binary fraction. First, some of the systems are more distant than others, making the tighter systems unresolvable. The importance of this effect is evident in Figure 4, which shows the magnitudes and colors of the target systems. Four of the ten systems with $I - K < 4$ and $K < 12$ are resolved as doubles, yet none of the seven systems with $I - K < 4$ and $K > 13.5$ are resolved. The latter are typically four times further away, making most double systems unresolvable. Figure 5 plots the histogram of the distance estimates. Overall, excluding the 16 systems in Figure 5 with photometric distances greater than 50 parsecs, *including the “wide” binary 2M1449+2355 and candidate binary 2M1600+1708*, the observed binary fraction is increased to $19 \pm 4\%$. Nevertheless, we may still fail to resolve many of the systems in the 1.6-10 A.U. range. This effect biases both the estimate of the binary frequency and the derived binary separation distribution. The observed binary separation distribution is plotted in Figure 6. In order to account for the systems which cannot be resolved, we also weight each system by the number of target systems in which we could have resolved it. For example, the binary 2M1017+1308 could have been resolved if it had been as close as 0.10

arcseconds; given our distance estimates it might have been resolved in only 57 of the total 102 targets searched. The resulting corrected distribution is also shown in Figure 6. It is evident that binaries in the range $\sim 2 - 5$ A.U. are more common than the wider systems.

A second important effect is that a luminosity-selected sample such as this one will be afflicted by systematic biases. Two conflicting biases exist (Reid 1991). First, near-equal luminosity binaries can be seen to greater distance and therefore are over-represented in the sample; if these extra doubles are resolved they will *increase* the observed binary fraction over the true one. Second, if these extra doubles are not resolved, they will *decrease* the observed binary fraction since they are counted as single stars. In the case of late-M and L dwarfs, where there are no observed systems with separations $\gtrsim 15$ A.U., the binaries may usefully be considered in two groups: Those with separations in the range $1.6 - 15$ A.U. and those with separation < 1.6 A.U. Our WFPC2 observations can resolve only the former group (with the innermost limit worse for the more distant objects), but through the second bias effect the latter group can affect our measurements.

The situation can be illustrated by a simple Monte Carlo model. For illustrative purposes only, we consider L dwarfs to be uniformly distributed in space out to a distance of 50 parsecs to a limiting magnitude of $K_s = 14.0$. In the present schematic model, we assign the same absolute magnitude, $M_K = 11.5$, to each primary. Secondary companions are added to a fraction f_1 of the L dwarfs at separations < 1 A.U. (unresolvable) and a fraction f_2 at $1 - 10$ A.U. (potentially resolvable). The K-band flux ratios are distributed uniformly in the range $0 < \Delta_K < 1$, and the f_2 dwarfs are given true separations between 1.0 and 10.0 . If the dwarf(s) have $K < 14$, they are considered as targets, and if they have apparent separation > 0.1 arcseconds they are considered to be resolved. We find that in models where the fraction of very-close binaries is very large (i.e., $f_1 \gtrsim 0.4$), then the WFPC2-resolved binary fraction is a significant underestimate of the true binary fraction f_2 . We therefore must consider what constraints exist on the fraction of very close, near-equal luminosity late-M and L dwarf binaries.

Dahn et al. (2002) have published trigonometric parallaxes of 2MASS and DENIS-selected L dwarfs. Of the 17 isolated field L dwarfs in their Table 1, and correcting for the five resolved companions in the range 1-10 A.U., only one (2M1328+21) appears to be offset from the L dwarf sequence and may be a close, unresolved double. This system was observed but not resolved by Reid et al. (2000). These data then suggest that $f_1 \lesssim 10\%$. This supported by the scarcity of double-lined binaries detected in high resolution studies of L dwarfs (Basri et al. 2000; Schweitzer et al. 2001). Similarly, Reid et al. (2002) find two double lined systems amongst a sample of 39 M6.5-L0.5 dwarfs. Other results come from studies of the Pleiades cluster. Martín et al. (2000b) estimate that 4 of 34 of Pleiades very-low-mass stars and brown dwarfs are potential near-equal luminosity binaries with separation < 27 A.U. based on their positions in the color-magnitude diagram. On the other hand, the double lined system PP1 15 (Basri & Martín 1999) shows close systems can exist.

The balance of evidence therefore suggests that $f_1 = 5 \pm 3\%$. It is then likely that the observed

binary percentage of $19 \pm 4\%$ is an overestimate of the true binary fraction (f_2) in the range 1.6-15 A.U. We have generated a series of Monte Carlo models similar to the ‘toy’ model described above, but using the observed distance distribution of the sample. By using this observed distance distribution, we include the most important factor in resolving binaries with HST. A more advanced treatment of the luminosity differences as a function of separation would be beyond the scope of this paper, and would depend on a better understanding of the sample selection effects, the substellar IMF, the L dwarf temperature scale, the star formation history, and other poorly known factors. Our best estimate, based on the simple Monte Carlo models, is that the true value $f_2 = 15 \pm 5\%$; the decrease due to the bias in including overluminous double is partially offset by the inability to resolve all binaries. On the other hand, the total binary fraction ($f_{tot} = f_1 + f_2$) must be larger than f_2 and is likely to be close to 20% under this model. Potential systematic error is comparable to the random errors due to sample size. In any case, it appears that a binary fraction $f_2 = 15 \pm 5\%$, with $15\% \leq f_{tot} \lesssim 25\%$. If enough L dwarfs within 10 parsecs can be discovered, HST and/or A.O. imaging may be able to improve the 1-3 A.U. constraints.

Since the binary fraction and observed orbital separations of the expanded sample are similar to those in the Reid et al. (2000) sample, the implications of the results remain the same. We refer the reader to Reid et al. (2000) and the more recent Close et al. (2003) for a discussion of the similarities and differences of the M and L dwarf binaries relative to main-sequence stars. In particular, the lack of systems at > 20 A.U. is now highly significant and a major difference from mid-M (Fischer & Marcy 1992) and G dwarfs (Duquennoy & Mayor 1991), as seen in Figure 7. A well-defined sample of M5-M6.5 dwarfs would allow the question of continuity of the binary frequency and orbital separation distribution to be addressed.

Since we cover both late-M, non-lithium L and lithium L dwarfs, we may search for differences as a function of mass of the primary. In other words, we address the question “does the binary fraction vary with mass near the hydrogen-burning limit?” Ten L dwarf systems in this study and Reid et al. (2000) have spectroscopic detections of lithium, placing them below the lithium-burning limit ($\sim 0.055M_\odot$).⁵ Three are doubles. Given the small sample size, the resulting binary fraction (30%) is consistent with that of the non-lithium L dwarfs and the late-M dwarfs. There is no evidence of a strong variation in binary frequency over the mass range ~ 0.08 to $\sim 0.04M_\odot$. It is interesting to note that all the doubles with companions in the range 10-16 A.U. have M8-L0 (probably stellar) dwarf primaries, while all of the later-L and lithium L (brown) dwarfs are tighter systems. Additional data are needed to assess the presently marginal significance of this effect. Our results are consistent with Burgasser et al. (2003)’s sample of ten T (brown) dwarfs in both the frequency and orbital separation. In particular, the fact that both of the resolved T dwarfs have separations < 5 A.U. is consistent with the peak seen for the late-M and L dwarfs separations in Figure 6.

⁵Three other L dwarf systems, all apparently single, have marginal lithium detections.

5. Summary and Conclusions

We have identified 13 definite and one candidate late-M and L dwarf binaries in a sample of 82 2MASS and SDSS selected field dwarfs. Including the twenty Reid et al. (2000) target systems, the observed binary fraction is $17^{+4}_{-3}\%$. We argue that accounting for biases and incompleteness leads to an estimated binary fraction $15 \pm 5\%$. No systems wider than 20 A.U. are found. Within the limits of the small sample sizes, we find no evidence for variations as a function of mass in the binary fraction. Our results are consistent with the (largely overlapping) late-M sample of Close et al. (2003) and the T dwarf sample of Burgasser et al. (2003).

Additional near-infrared observations of the brown dwarf double 2M1728+3948 are needed. We interpret the secondary as an early-T dwarf. Orbital motions for these systems should be detectable using HST. This offers the opportunity to determine masses.

The greatest need for the future is a better-defined sample of L dwarf targets, such as a 2MASS-selected sample of bright L dwarfs (Cruz et al., in prep.). The current sample of L dwarfs, although large, was not selected with consistent magnitude and color cuts that have well-understood effects. Once such a sample is available with HST or AO data, the bias and selection effects can be better understood.

Based on observations made with the NASA/ESA Hubble Space Telescope, obtained at the Space Telescope Science Institute, which is operated by the Association of Universities for Research in Astronomy, Inc., under NASA contract NAS 5-26555. These observations are associated with proposal 8581. Support for this work was provided by NASA through grant number HST-GO-08581.01-A from the Space Telescope Science Institute, which is operated by AURA, Inc., under NASA contract NAS 5-26555. This publication makes use of data products from 2MASS, which is a joint project of the University of Massachusetts and IPAC/Caltech, funded by NASA and NSF.

REFERENCES

- Basri, G., & Martín, E.L. 1999, *AJ*, 118, 2460
- Basri, G., et al. 2000, *ApJ*, 538, 363
- Burgasser, A.J., et al. 2002a, *ApJ*, 564, 421
- Burgasser, A.J., et al. 2002b, *ApJ*, 571, L151
- Burgasser, A. J., Reid, I. N., Kirkpatrick, J. D., Brown, M. E., Miskey, C. L., & Gizis, J. E. 2003, *ApJ*, submitted
- Burrows, A., Hubbard, W.B., Lunine, J.I., & Liebert, J. 2001, *Rev. Modern Physics*, 73, 719
- Chabrier, G., & Baraffe, I. 1997, *A&A*, 327, 1039
- Close, L.M., Potter, D., Brandner, W., Lloyd-Hart, M., Liebert, J., Burrows, A., & Siegler, N. 2002a, *ApJ*, 566, 1095
- Close, L.M., Siegler, N., Potter, D., Brandner, W., & Liebert, J. 2002b, *ApJ*, 567, L53
- Close, L.M., Siegler, N., Freed, M., & Biller, B. 2003, *ApJ*, in press (astro-ph/0301095)
- Cutri, R.M., et al. 2000, Explanatory Supplement to the 2MASS Second Incremental Data Release, <http://www.ipac.caltech.edu/2mass/releases/second/doc/explsup.html>
- Dahn, C.C., et al. 2002, *AJ*, 124, 1170
- Delfosse, X., et al. 1997, *A&A*, 327, L25
- Dolphin, A.E. 2000, *PASP*, 112, 1397
- Duquennoy, A. & Mayor, M. 1991, *A&A*, 248, 485
- Epchtein, N. 1997, in *The Impact of Large Scale Near-IR Sky Surveys*, eds. F. Garzon et al., p. 15
- Fan, X., et al. 2000, *AJ*, 119, 928
- Fischer, D.A., & Marcy, G.W. 1992, *ApJ*, 396, 178
- Geballe, T.R., et al. 2002, *ApJ*, 564, 466
- Gizis, J.E., & Reid, I.N. 1995, *AJ*, 110, 1248
- Gizis, J.E., Monet, D.G., Reid, I.N., Kirkpatrick, J.D., Liebert, J. & Williams, R.J. 2000, *AJ*, 120, 1085
- Halbwachs, J. L., Arenou, F., Mayor, M., Udry, S., & Queloz, D. 2000, *A&A*, 355, 581

- Hawley, S.L., et al. 2002, *AJ*, 123, 3409
- Holtzman, J.A., Burrows, C.J., Casertano, S., Hester, J.J., Trauger, J.T., Watson, A.M., & Worthey, G. 1995, *PASP*, 107, 1065
- Joyce, R. 1999, The CRSP User’s Manual, <http://www.noao.edu/kpno/manuals/crsp/>
- Kirkpatrick, J.D., Reid, I.N., Liebert, J., Cutri, R.M., Nelson, B., Beichman, C.A., Dahn, C.C., Monet, D.G., Gizis, J.E., & Skrutskie, M.F. 1999, *ApJ*, 519, 802
- Kirkpatrick, J.D., Reid, I.N., Gizis, J.E., Burgasser, A.J., Liebert, J., Monet, D.G., Dahn, C.C., Nelson, B. 2000, *AJ*, 120, 447
- Koerner, D.W., Kirkpatrick, J.D., McElwain, M.W., & Bonaventura, N.R. 1999, *ApJ*, 526, L25
- Martín, E.L., Delfosse, X., Basri, G., Goldman, B., Forveille, T., Zapatero Osorio, M.R. 1999a, *AJ*, 118, 2466
- Martín, E. L., Brandner, W., & Basri, G. 1999b, *Science*, 283, 1718
- Martín, E. L., Koresko, C. D., Kulkarni, S. R., Lane, B. F., & Wizinowich, P. L. 2000a, *ApJ*, 529, L37
- Martín, E. L., et al. 2000b, *ApJ*, 543, 299
- Marcy, G.W., & Butler, R.P. 2000, *PASP*, 112, 137
- Potter, D., Martín, E. L., Cushing, M. C., Baudoz, P., Brandner, W., Guyon, O., & Neuhauser, R. 2002, *ApJ*, 567, L133
- Reid, N. 1991, *AJ*, 102, 1428
- Reid, I.N., Kirkpatrick, J.D., Liebert, J., Burrows, A., Gizis, J.E., Burgasser, A., Dahn, C.C., Monet, D., Cutri, R., Beichman, C.A., & Skrutskie, M. 1999, *ApJ*, 521, 613
- Reid, I.N., Gizis, J.E., Kirkpatrick, J.D., & Koerner, D.W. 2000, *AJ*, 121, 489
- Reid, I.N., Kirkpatrick, J.D., Liebert, J., Gizis, J.E., Dahn, C.C., & Monet, D.G. 2002, *AJ*, 124, 519
- Schneider, D.P., et al. 2002, *AJ*, 123, 458
- Schweitzer, A., Gizis, J.E., Hauschildt, P.H., Allard, F., & Reid, I.N. 2001, *ApJ*, 555, 363
- Schweitzer, A., Gizis, J.E., Hauschildt, P.H., Allard, F., Howard, E.M., Kirkpatrick, J. D. 2002, *ApJ*, 566, 435

Table 1. Target Systems

Name	I	Z	J	H	K_s	σ_I	σ_Z	σ_J	σ_H	σ_K	d (pc)	No.	Ref.
2MASS J03264225-2102057	19.977	17.422	16.111	14.774	13.885	0.027	0.091	0.100	0.075	0.062	14	1	i
2MASS J08564793+2235182	19.096	16.888	15.647	14.579	13.924	0.041	0.039	0.061	0.055	0.050	35	1	h
2MASS J00100368+3436099	18.231	16.332	15.642	15.080	14.391	0.026	0.059	0.066	0.089	0.073	97	1	e
2MASS J00283943+1501418	20.439	17.970	16.477	15.226	14.539	0.042	0.127	0.109	0.086	0.071	12	1	f
2MASS J00303013-1450333	20.286	17.752	16.325	15.283	14.492	0.029	0.101	0.115	0.102	0.097	11	1	f
2MASS J00332386-1521309	18.718	16.612	15.294	14.225	13.397	0.020	0.058	0.055	0.043	0.040	31	1	i
2MASS J02085499+2500488	17.196	15.083	14.015	13.110	12.579	0.017	0.031	0.032	0.035	0.037	27	1	f
2MASS J02243670+2537042	19.760	17.722	16.550	15.419	14.670	0.034	0.086	0.107	0.083	0.085	83	1	f
2MASS J03284265+2302051	20.434	17.493	16.556	15.547	14.833	0.031	0.111	0.155	0.132	0.105	17	1	f
2MASS J03350208+2342356	14.687	...	12.259	11.654	11.261	0.021	...	0.028	0.033	0.025	21	1	c
2MASS J03370359-1758079	19.405	17.019	15.594	14.412	13.588	0.021	0.058	0.065	0.051	0.047	13	1	f
2MASS J03454316+2540233	17.244	14.996	13.992	13.170	12.665	0.034	0.020	0.028	0.034	0.029	27.0	1	
2MASS J03505737+1818069	15.603	13.733	12.951	12.222	11.763	0.017	0.022	0.031	0.034	0.032	28	1	c
2MASS J03554191+2257016	19.545	17.318	16.099	15.023	14.247	0.024	0.064	0.087	0.075	0.066	43	1	e
2MASS J07533217+2917119	18.802	16.595	15.485	14.489	13.849	0.022	0.046	0.049	0.047	0.055	42	1	f
2MASS J08014056+4628498	19.757	17.356	16.287	15.439	14.540	0.029	0.069	0.140	0.146	0.112	45	1	f
2MASS J08295707+2655099	20.774	18.238	17.109	15.723	14.685	0.028	0.113	0.201	0.140	0.081	40	1	f
2MASS J08320451-0128360	17.388	15.283	14.127	13.309	12.687	0.023	0.026	0.028	0.027	0.031	25	1	f
2MASS J09141884+2238134	18.391	16.280	15.304	14.396	13.898	0.021	0.036	0.047	0.045	0.041	56	1	e
2MASS J09510549+3558021	20.971	18.177	17.358	15.894	14.966	0.026	0.143	0.256	0.155	0.101	52	1	f
2MASS J10170754+1308398	17.414	15.275	14.126	13.194	12.683	0.040	0.047	0.033	0.034	0.032	30	2	m
2MASS J11023375-2359464	20.812	18.398	17.040	15.614	14.794	0.025	0.114	0.193	0.125	0.112	29	1	f
2MASS J11040127+1959217	17.981	15.751	14.369	13.492	12.974	0.019	0.028	0.031	0.038	0.038	13	1	h
2MASS J11083081+6830169	16.554	-7.037	13.139	12.227	11.600	0.027	0.000	0.027	0.024	0.031	12	1	c
2MASS J11122567+3548131	18.144	15.990	14.573	13.473	12.694	0.064	0.075	0.043	0.041	0.046	16	1	f
2MASS J11275346+7411076	15.812	13.871	13.059	12.367	11.971	0.041	0.032	0.031	0.028	0.029	36	2	c
2MASS J12391934+2029519	17.326	15.225	14.465	13.611	13.116	0.018	0.035	0.034	0.038	0.036	49	1	e
2MASS J12392727+5515371	18.338	16.009	14.670	13.539	12.743	0.032	0.060	0.033	0.036	0.031	19	2	f
2MASS J13113921+8032219	15.458	13.552	12.784	12.116	11.721	0.039	0.037	0.022	0.025	0.024	33	2	c
2MASS J14032232+3007547	15.415	13.482	12.691	12.008	11.626	0.024	0.027	0.026	0.028	0.026	24	1	c

Table 1—Continued

Name	<i>I</i>	<i>Z</i>	<i>J</i>	<i>H</i>	<i>K_s</i>	σ_I	σ_Z	σ_J	σ_H	σ_K	d (pc)	No.	Ref.
2MASS J17433487+5844110	17.123	14.762	14.016	13.153	12.669	0.037	0.095	0.025	0.030	0.031	30	1	h
2MASS J18410861+3117279	19.746	17.393	16.120	14.970	14.180	0.028	0.056	0.100	0.070	0.084	28	1	f
2MASS J20543585+1519043	19.660	17.485	16.205	15.443	14.811	0.031	0.059	0.107	0.128	0.110	44	1	f
2MASS J20571538+1715154	19.357	17.268	16.107	15.211	14.567	0.031	0.082	0.110	0.090	0.128	63	1	f
2MASS J21011544+1756586	20.677	18.155	16.825	15.792	15.173	0.043	0.188	0.178	0.157	0.194	26	2	f
2MASS J21402931+1625183	15.723	13.764	12.943	12.270	11.779	0.029	0.027	0.031	0.034	0.033	32	2	c
2MASS J21474365+1431315	16.857	14.705	13.842	13.134	12.652	0.051	0.093	0.036	0.032	0.042	40	2	c
2MASS J21580457-1550098	18.559	16.263	14.949	13.916	13.148	0.024	0.060	0.037	0.043	0.040	17	1	l
2MASS J22062280-2047058	15.007	13.197	12.381	11.704	11.325	0.031	0.034	0.026	0.023	0.029	31	2	c
2MASS J22064500-4217210	19.143	16.913	15.569	14.478	13.595	0.028	0.053	0.068	0.056	0.057	25	1	f
2MASS J22081363+2921215	19.588	...	15.818	14.825	14.086	0.028	0.427	0.090	0.083	0.083	17	1	f
2MASS J22244381-0158521	17.759	15.363	14.052	12.803	12.017	0.016	0.034	0.030	0.029	0.029	11.4	1	f
2MASS J22341394+2359559	16.247	14.156	13.176	12.353	11.835	0.031	0.022	0.022	0.030	0.034	22	1	c
2MASS J22425317+2542573	18.124	15.977	14.795	13.754	13.022	0.025	0.031	0.038	0.036	0.038	30	1	i
2MASS J22443167+2043433	20.393	17.678	16.405	14.965	13.932	0.026	0.067	0.128	0.071	0.066	11	1	a
2MASS J23062928-0502285	14.004	...	11.372	10.718	10.288	0.013	...	0.022	0.031	0.027	13	1	c
2MASS J23094618+1549045	17.789	15.847	15.005	14.343	13.907	0.034	0.039	0.059	0.058	0.065	67	1	e
2MASS J23310161-0406193	15.536	13.722	12.937	12.289	11.930	0.027	0.030	0.027	0.027	0.029	28	2	c
2MASS J23494899+1224386	15.320	13.449	12.615	11.952	11.562	0.023	0.022	0.024	0.026	0.030	23	1	c
SDSS J001911.65+003017.8	18.174	15.991	14.924	14.173	13.588	0.020	0.044	0.036	0.034	0.037	37	1	d
2MASS J03440892+0111251*	17.872	15.800	14.725	13.890	13.522	0.017	0.034	0.035	0.036	0.047	40	1	k
SDSSp J104325.10+000148.2	19.255	17.041	16.096	15.111	14.709	0.027	0.059	0.098	0.082	0.120	73	1	g
SDSS J143535.72-004347.0	19.747	17.653	16.461	15.672	15.073	0.022	0.088	0.111	0.115	0.141	70	1	d
SDSS J143517.20-004612.9	19.142	17.202	16.444	15.676	15.268	0.032	0.068	0.104	0.118	0.174	136	1	d
2MASS J15483164-0029414*	18.230	16.532	15.689	15.123	14.850	0.060	0.035	0.067	0.083	0.125	100	1	k
SDSS J165329.69+623136.5	18.371	16.280	15.109	14.389	13.864	0.026	0.033	0.054	0.053	0.067	39	1	d
2MASS J17232861+6406230*	18.961	17.046	16.323	15.644	15.175	0.018	0.051	0.111	0.142	0.191	131	1	k
2MASS J23355849-0013042*	18.735	16.849	15.969	15.206	14.688	0.058	0.091	0.079	0.089	0.104	105	1	k
SDSSp J033035.13-002534.5	18.855	16.520	15.290	14.419	13.829	0.019	0.049	0.045	0.042	0.051	23	1	b
SDSSp J053951.99-005902.0	17.575	15.228	13.986	13.065	12.577	0.019	0.026	0.028	0.028	0.030	12	1	b
2MASS J15154719-0030594*	16.985	15.034	14.181	13.578	13.144	0.016	0.032	0.027	0.028	0.038	45	1	b

Skrutskie, M.F. et al. on The Impact of Large Scale Near-IR Sky Surveys, eds. F. Garzon et al., p. 25.

York, D.G., et al. 2000, AJ, 120, 1579

Table 2. Doubles

Name	I_A	Z_A	σ_{I_A}	σ_{Z_A}	I_B	Z_B	σ_{I_B}	σ_{Z_B}	Sep. (")	PA	Sep. (AU)
2M1017+1308	18.080	15.905	0.035	0.043	18.260	16.165	0.034	0.045	89	0.100	3
2M1127+7411	16.370	14.478	0.036	0.025	16.803	14.792	0.021	0.023	81	0.250	9
2M1239+5515	19.077	16.675	0.025	0.057	19.104	16.856	0.025	0.055	7	0.213	4
2M1311+8032	16.105	14.223	0.033	0.031	16.328	14.393	0.029	0.024	167	0.300	10
2M1426+1557	16.148	14.108	0.034	0.035	17.291	15.218	0.026	0.023	340	0.155	4
2M1430+2915	18.106	15.947	0.036	0.036	18.628	16.422	0.049	0.048	326	0.084	3
2M1449+2355	19.115	16.984	0.018	0.080	20.151	17.829	0.027	0.113	64	0.133	13
2M1600+1708 ^a	19.912	17.644	0.048	0.157	20.072	17.942	343	0.101	0.218	0.056	6
2M1728+3948	20.268	17.802	0.029	0.084	20.636	17.551	0.028	0.110	27	0.130	3
2M2101+1756	21.226	18.579	0.038	0.187	21.681	19.381	0.045	0.343	107	0.232	6
2M2140+1625	16.030	14.130	0.021	0.018	17.245	15.122	0.031	0.043	132	0.158	5
2M2147+1431	17.430	15.305	0.047	0.091	17.825	15.636	0.036	0.048	329	0.323	13
2M2206-2047	15.721	13.940	0.024	0.027	15.799	13.959	0.024	0.024	57	0.160	5
2M2331-0406	15.568	13.784	0.018	0.022	19.363	16.864	0.039	0.066	294	0.576	16

^aCandidate binary




Fig. 1.— KPNO 4-meter CRSP spectra of three new L dwarfs. The observations are described in Schweitzer et al. (2002).




Fig. 2.— Color-color diagram using HST (I,Z) and 2MASS (J) photometry. Apparently single dwarfs are shown as crosses and doubles as open squares. T dwarfs from Burgasser et al. (2003) are shown as solid points.




Fig. 3.— Color-color diagram using HST (I,Z) and 2MASS (J) photometry. Symbols as in Figure 2




Fig. 4.— Color-magnitude diagram for the sample without corrections for binarity. The apparently single systems appear as crosses and the resolved doubles as open squares. The solid curves plot the Dahn et al. (2002) cool dwarf sequence at distances of 10, 25, and 50 parsecs.




Fig. 5.— Histogram of estimated photometric parallaxes.




Fig. 6.— Histogram of observed orbital separations. The histogram shows the count of resolved systems, while the solid points with connecting lines shows the distribution after correction for the fact that the closest systems cannot be resolved in all cases. The distribution is peaked, with companions more common in the range 2-5 A.U. in the range 6-10 A.U.




Fig. 7.— Derived binary fractions for late-M and L dwarfs compared to early-M dwarfs (long-dashed lines, Fischer & Marcy 1992) and G dwarfs (short-dashed lines, Duquennoy & Mayor 1991); all stellar companions regardless of mass ratio are included. Note the lack of wide companions and the relative “excess” of 1-16 A.U. companions for the coolest primaries relative to the G dwarfs.